

急性曝露ガイドライン濃度 (AEGL)

Potassium Phosphide (20770-41-6)

リン化カリウム

Table AEGL 設定値

Potassium Phosphide 20770-41-6 (Final)					
ppm					
	10 min	30 min	60 min	4 hr	8 hr
AEGL 1	NR	NR	NR	NR	NR
AEGL 2	4.0	4.0	2.0	0.50	0.25
AEGL 3	7.2	7.2	3.6	0.90	0.45

NR: データ不十分により推奨濃度設定不可

設定根拠 (要約) :

ホスフィンは無色の気体で、貯蔵穀物を食害する昆虫やげっ歯類に対し燻蒸剤として使用される。駆除剤は、金属リン化物の形で使用されることが一般的であり、水分と反応してホスフィンガスを遊離する。ホスフィンも、半導体産業で使用される。ホスフィンのヒトへの曝露に関するデータは、曝露の時間と濃度が正確に報告されていないため、AEGL値の導出に使用するには制限がある。一方、動物については、適切なデータは豊富にあるものの、AEGL-1値の定義に合致するデータは得られていない。したがって、AEGL-1値は、データ不十分のため、導出しなかった。

AEGL-2値は、Newtonら (1993) の試験で、10 ppmのホスフィンに6時間曝露したFischer 344 ラットにおいて、赤色の粘液性鼻汁が認められたことに基づいた。ラット、マウス、ウサギ、およびモルモットの45分間～30時間の致死データ (死亡までの時間) から、種差はほとんどないことが示唆されるため (Figure 10-2を参照)、種間変動に関する不確実係数として3を適用した。ホスフィンへの曝露濃度が同じであると推定される場合、小児は成人よりもホスフィンに対する感受性が高いことが、ヒトにおけるデータから示唆されるため、種内変動に関する不確実係数として10を適用した。したがって、総不確実係数は30となる。全身に作用する刺激性の蒸気やガスの多くは、曝露濃度-曝露時間関係を $C^n \times t = k$ の式で表すことができ、指数 n は0.8～3.5の範囲の値をとる (ten Berge et al. 1986)。ホスフィンについて、30分間、1時間、4時間、8時間の各曝露時間のAEGL値をスケールリングするため、指数 n には、経験に基づいて導出した1の値を適用した。また、指数 n の導出には、ラットにお

ける1～6時間の致死データを使用した。6時間から10分間への外挿に係る不確実性が加わるため、AEGL-2の30分間値を10分間値としても採用した。




FIGURE 10-2 Best-fit concentration (ppm phosphine) × time (exposure duration in hours) curve. Linear regression of rat lethality data.

AEGL-3値は、ホスフィンに6時間曝露したSprague Dawleyラットにおいて、死亡が認められなかった濃度（18 ppm）に基づいた。種差はほとんどないことが、ラット、マウス、ウサギ、モルモットの致死データから示唆されるため、種間変動に関する不確実係数として3を適用した。ホスフィンの曝露濃度が同じであると推定される場合、小児は成人よりもホスフィンに対する感受性が高いことが、ヒトにおけるデータから示唆されるため、種内変動に関する不確実係数として10を適用した。したがって、総不確実係数は30となる。全身に作用する刺激性の蒸気やガスの多くは、曝露濃度-曝露時間関係を $C^n \times t = k$ の式で表すことができ、指数 n は0.8～3.5の範囲の値をとる (ten Berge et al. 1986)。ホスフィンについては、30分間、1時間、4時間、8時間の各曝露時間のAEGL値をスケールリングするため、指数 n には、経験に基づいて導出した1の値を適用した。指数 n の導出には、ラットにおける1～6時間の致死データを使用した。6時間から10分間への外挿に係る不確実性が加わるため、AEGL-3の30分間値を10分間値としても採用した。Table に、導出したAEGL値を示す。

Appendix D に、8種類の金属リン化物のAEGL値を示す。

APPENDIX D**AEGL Values for Selected Metal Phosphides****Aluminum Phosphide (AlP)****Potassium Phosphide (K₃P)****Sodium Phosphide (Na₃P)****Zinc Phosphide (Zn₃P₂)****Calcium Phosphide (Ca₃P₂)****Magnesium Phosphide (Mg₃P₂)****Strontium Phosphide (Sr₃P₂)****Magnesium Aluminum Phosphide (Mg₃AlP₃)****SUMMARY**

Metal phosphides are solids and are typically used as fumigants against insects and rodents in stored grain. The metal phosphides react rapidly with water and moisture in the air or stored grain to produce phosphine gas. It is the phosphine gas that is responsible for acute toxicity, and the rate of phosphine generation is dependent on ambient temperature and humidity and the chemical structure of the phosphide (Anger et al. 2000).

In the absence of appropriate chemical-specific data for the metal phosphides considered in this appendix, the AEGL-2 and AEGL-3 values for phosphine were used to obtain AEGL-2 and AEGL-3 values, respectively, for the metal phosphides. The use of phosphine as a surrogate for the metal phosphides is deemed appropriate because qualitative (clinical signs) and quantitative (phosphine blood level) data suggest that the phosphine hydrolysis product is responsible for acute toxicity from metal phosphides. The phosphine AEGL-2 and AEGL-3 values were used as target values for calculating the concentrations of metal phosphide needed to generate the phosphine AEGL values.

Because AEGL-1 values for phosphine are not recommended (due to insufficient data), AEGL-1 values for the metal phosphides considered in this appendix are also not recommended. The calculated values are listed in Table D-1 below.

D.I. INTRODUCTION



Metal phosphides are solids and are typically used as fumigants against insects and rodents in stored grain. The metal phosphides react rapidly with water and moisture in the air or stored grain to produce phosphine gas. It is the phosphine gas which is responsible for acute toxicity, and the rate of phosphine generation is dependent on ambient temperature and humidity and the chemical structure of the phosphide (Anger et al. 2000).

TABLE D-1 AEGL Values for Metal Phosphides^a





Compound	Classification	10-min	30-min	1-hr	4-hr	8-hr
Aluminum Phosphide	AEGL-1	NR	NR	NR	NR	NR
	AEGL-2	9.5 mg/m ³	9.5 mg/m ³	4.7 mg/m ³	1.2 mg/m ³	0.59 mg/m ³
	AEGL-3	17 mg/m ³	17 mg/m ³	8.5 mg/m ³	2.1 mg/m ³	1.1 mg/m ³
Potassium Phosphide	AEGL-1	NR	NR	NR	NR	NR
	AEGL-2	24 mg/m ³	24 mg/m ³	12 mg/m ³	3.0 mg/m ³	1.5 mg/m ³
	AEGL-3	44 mg/m ³	44 mg/m ³	22 mg/m ³	5.5 mg/m ³	2.7 mg/m ³
Sodium Phosphide	AEGL-1	NR	NR	NR	NR	NR
	AEGL-2	16 mg/m ³	16 mg/m ³	8.2 mg/m ³	2.0 mg/m ³	1.0 mg/m ³
	AEGL-3	29 mg/m ³	29 mg/m ³	15 mg/m ³	3.7 mg/m ³	1.8 mg/m ³
Zinc Phosphide	AEGL-1	NR	NR	NR	NR	NR
	AEGL-2	21 mg/m ³	21 mg/m ³	11 mg/m ³	2.6 mg/m ³	1.3 mg/m ³
	AEGL-3	38 mg/m ³	38 mg/m ³	19 mg/m ³	4.8 mg/m ³	2.4 mg/m ³
Calcium Phosphide	AEGL-1	NR	NR	NR	NR	NR
	AEGL-2	15 mg/m ³	15 mg/m ³	7.5 mg/m ³	1.9 mg/m ³	0.93 mg/m ³
	AEGL-3	27 mg/m ³	27 mg/m ³	13 mg/m ³	3.4 mg/m ³	1.7 mg/m ³
Magnesium Phosphide	AEGL-1	NR	NR	NR	NR	NR
	AEGL-2	11 mg/m ³	11 mg/m ³	5.5 mg/m ³	1.4 mg/m ³	0.69 mg/m ³
	AEGL-3	20 mg/m ³	20 mg/m ³	9.9 mg/m ³	2.5 mg/m ³	1.2 mg/m ³
Strontium Phosphide	AEGL-1	NR	NR	NR	NR	NR
	AEGL-2	27 mg/m ³	27 mg/m ³	13 mg/m ³	3.3 mg/m ³	1.7 mg/m ³
	AEGL-3	48 mg/m ³	48 mg/m ³	24 mg/m ³	6.0 mg/m ³	3.0 mg/m ³
Magnesium Aluminum Phosphide	AEGL-1	NR	NR	NR	NR	NR
	AEGL-2	11 mg/m ³	11 mg/m ³	5.3 mg/m ³	1.3 mg/m ³	0.66 mg/m ³
	AEGL-3	19 mg/m ³	19 mg/m ³	9.5 mg/m ³	2.4 mg/m ³	1.2 mg/m ³

^aThese airborne concentrations will produce the equivalent AEGL values for phosphine.
 Note: Absence of an AEGL-1 does not imply that exposure below the AEGL-2 is without adverse effects.
 NR, not recommended.

Aluminum Phosphide (CAS No. 20859-73-8), Potassium Phosphide (CAS No. 20770-41-6), and Sodium Phosphide (CAS No. 12058-85-4): One mole of aluminum phosphide, potassium phosphide, or sodium phosphide will react rapidly with water or moisture to produce a maximum of 1 mole of phosphine gas as follows:



Zinc Phosphide (CAS No. 1314-84-7), Calcium Phosphide (CAS No. 1305-99-3), Magnesium Phosphide (CAS No. 10257-74-8), and Strontium Phosphide (CAS No. 12504-13-1): One mole of zinc phosphide, calcium phosphide, magnesium phosphide or strontium phosphide will react rapidly with water or moisture to produce a maximum of 2 moles of phosphine gas as follows:



Magnesium Aluminum Phosphide (CAS No. None): One mole of magnesium aluminum phosphide will react rapidly with water or moisture to produce a maximum of 3 moles of phosphine gas as follows:



Aluminum phosphide is a gray or yellow crystalline solid prepared from red phosphorus and aluminum powder (O'Neil et al. 2001). Commercial aluminum phosphide sachets contain 70% aluminum phosphide and 30% aluminum carbonate (Bajaj et al. 1988). Calcium phosphide is a red-brown or gray solid prepared by heating calcium phosphate with aluminum or carbon by passing phosphorus vapors over metallic calcium. In addition to its use as a rodenticide, calcium phosphide is also used in signal fires and pyrotechnics and in the purification of copper and copper alloys (HSDB 2007a). Zinc phosphide is a gray solid and may be produced by passing phosphine through a solution of zinc sulfate (HSDB 2007b). Manufacturing information on the other metal phosphides considered in this appendix was not located. Available physico-chemical data for the metal phosphides are shown in Tables D-2 through D-9.

TABLE D-2 Physicochemical Data for Aluminum Phosphide

Parameter	Description/Value	Reference
Synonyms (commercial product)	Celphos, Phostoxin, Quickphos	O'Neil et al. 2001
CAS Registry No.	20859-73-8	O'Neil et al. 2001
Chemical formula	AlP	O'Neil et al. 2001
Molecular weight	57.96	O'Neil et al. 2001
Physical state	Solid, gray of yellow crystals	O'Neil et al. 2001

(Continued)

TABLE D-2 Continued

Parameter	Description/Value	Reference
Relative density (water = 1)	2.9	IPCS 1989
Melting point	>1350°C	IPCS 1989
Solubility in water	Reactive produces phosphine gas	IPCS 1989

TABLE D-3 Physicochemical Data for Potassium Phosphide

Parameter	Description/Value	Reference
CAS Registry No.	20770-41-6	ChemIDPlus 2005a
Chemical formula	K ₃ P	ChemIDPlus 2005a
Molecular weight	148.3	ChemIDPlus 2005a

TABLE D-4 Physicochemical Data for Sodium Phosphide

Parameter	Description/Value	Reference
Synonyms	Trisodium phosphide	ChemIDPlus 2005b
CAS Registry No.	12058-85-4	ChemIDPlus 2005b
Chemical formula	Na ₃ P	ChemIDPlus 2005b
Molecular weight	99.94	Lewis 1996a
Physical state	Solid, red crystals	Lewis 1996a
Melting point	Decomposes	Lewis 1996a
Solubility in water	Reacts violently	Lewis 1996a

TABLE D-5 Physicochemical Data for Zinc Phosphide

Parameter	Description/Value	Reference
Synonym	Trizinc diphosphide	IPCS 1989
CAS Registry No.	1314-84-7	IPCS 1989
Chemical formula	Zn ₃ P ₂	IPCS 1989
Molecular weight	258.1	IPCS 1989
Physical state	Solid, gray powder or crystals	O'Neil et al. 2001
Relative density (water = 1)	4.55	O'Neil et al. 2001
Melting point	Sublimes	IPCS 1989
Solubility in water	Insoluble, reacts	IPCS 1989

TABLE D-6 Physicochemical Data for Calcium Phosphide

Parameter	Description/Value	Reference
Synonyms (commercial product)	Calcium photophor, photophor	O'Neil et al. 2001
CAS Registry No.	1305-99-3	O'Neil et al. 2001
Chemical formula	Ca ₃ P ₂	O'Neil et al. 2001

(Continued)

TABLE D-6 Continued

Parameter	Description/Value	Reference
Molecular weight	182.18	O'Neil et al. 2001
Physical state	Solid, red-brown crystals or gray lumps	O'Neil et al. 2001
Relative density (water = 1)	2.51	O'Neil et al. 2001
Melting point	1600°C	O'Neil et al. 2001
Solubility in water	Decomposes	O'Neil et al. 2001

TABLE D-7 Physicochemical Data for Magnesium Phosphide

Parameter	Description/Value	Reference
Synonyms	Trimagnesium diphosphide	IPCS 1989
CAS Registry No.	12057-74-8	IPCS 1989
Chemical formula	Mg ₃ P ₂	IPCS 1989
Molecular weight	134.87	IPCS 1989
Physical state	Solid, gray or bright yellow crystals	IPCS 1989
Relative density (water = 1)	2.1	IPCS 1989
Melting point	>750°C	IPCS 1989
Solubility in water	Moisture sensitive, reacts	IPCS 1989

TABLE D-8 Physicochemical Data for Strontium Phosphide

Parameter	Description/Value	Reference
CAS Registry No.	12504-13-1	Lewis 1996b
Chemical formula	Sr ₃ P ₂	Lewis 1996b
Molecular weight	324.9	Lewis 1996b
Physical state	Solid	Lewis 1996b

TABLE D-9 Physicochemical Data for Magnesium Aluminum Phosphide

Parameter	Description/Value	Reference
CAS Registry No.	None	ChemIDPlus 2005c
Chemical formula	Mg ₃ AlP ₃	ChemIDPlus 2005c
Molecular weight	192.8	—
Physical state	Solid	ChemIDPlus 2005c

D.II. SPECIAL CONSIDERATIONS

Metabolism and Disposition

Solid metal phosphides deposited on moist respiratory tract surfaces may hydrolyze and release absorbable phosphine. However, a more likely scenario would involve atmospheric hydrolysis of metal phosphides to phosphine gas.

Chan et al. (1983) detected phosphine in postmortem stomach, blood, and liver specimens from a 27-year-old man who had ingested aluminum phosphide tablets. The phosphine was released from the samples after acid treatment. Similarly, Anger et al. (2000) identified phosphine in postmortem brain, liver, and kidneys of a 39-year-old man who had committed suicide by ingestion of aluminum phosphide tablets.

Chugh et al. (1996) measured blood phosphine levels in patients with severe ($n = 30$), mild ($n = 10$), or minimal ($n = 5$) toxicity due to aluminum phosphide ingestion. Patients with severe toxicity had ingested “fresh” aluminum phosphide compound and were in a state of shock. Those with mild toxicity had ingested “old” aluminum phosphide compound and presented with hypotension and gastrointestinal symptoms. Patients with minimal toxicity ingested some powder from the aluminum phosphide tablets and presented with only nausea and occasional vomiting. Blood phosphine levels were positively correlated with severity of clinical signs and to dose of pesticide. At admission, blood phosphine levels were 71% higher ($p < 0.001$) in patients in the severe toxicity group than in the mild toxicity group of patients; blood phosphine was not detected in the minimal toxicity group of patients. Blood phosphine levels were also correlated to mortality; patients having blood phosphine levels $\leq 1.067 \pm 0.16$ mg% survived, whereas those with blood phosphine above this apparent threshold died (6 of 30 in the severe toxicity group).

Garry et al. (1993) described a fatality from inhalation of aluminum phosphide aerosol. In this case report, blood aluminum concentration was used as a marker of exposure (see Section 2.1.1).

Mechanism of Toxicity

Metal phosphides react rapidly with moisture in air to produce phosphine gas. It is the phosphine gas that is responsible for acute inhalation toxicity from metal phosphide exposure. The rate of phosphine generation is dependent on ambient temperature and humidity (Anger et al. 2000) in addition to the chemical structure of the metal phosphide. The hydrolysis reactions and phosphine evolution rates (OECD 2001) of the metal phosphides considered in this appendix are summarized in Table D-10.

D.III. RATIONALE AND AEGL-1

Summary of Human Data Relevant to AEGL-1

No human data are available for the derivation of AEGL-1 for the metal phosphides considered in this appendix.

TABLE D-10 Hydrolysis of Metal Phosphides

Metal Phosphide	Hydrolysis Reaction	Maximum Number of moles of phosphine produced per mole of metal phosphide hydrolyzed	Phosphine evolution rate at 20°C and 1 atm (mL/kg•min)
Aluminum Phosphide	$\text{AlP} + 3\text{H}_2\text{O} \rightarrow \text{PH}_3 + \text{Al}(\text{OH})_3$	1	2069.7
Potassium Phosphide	$\text{K}_3\text{P} + 3\text{H}_2\text{O} \rightarrow \text{PH}_3 + 3\text{KOH}$	1	807.6
Sodium Phosphide	$\text{Na}_3\text{P} + 3\text{H}_2\text{O} \rightarrow \text{PH}_3 + 3\text{NaOH}$	61	997.8
Zinc Phosphide	$\text{Zn}_3\text{P}_2 + 6\text{H}_2\text{O} \rightarrow 2\text{PH}_3 + 3\text{Zn}(\text{OH})_2$	2	929.9
Calcium Phosphide	$\text{Ca}_3\text{P}_2 + 6\text{H}_2\text{O} \rightarrow 2\text{PH}_3 + 3\text{Ca}(\text{OH})_2$	2	1274.6
Magnesium Phosphide	$\text{Mg}_3\text{P}_2 + 6\text{H}_2\text{O} \rightarrow 2\text{PH}_3 + 3\text{Mg}(\text{OH})_2$	2	1781.4
Strontium Phosphide	$\text{Sr}_3\text{P}_2 + 6\text{H}_2\text{O} \rightarrow 2\text{PH}_3 + 3\text{Sr}(\text{OH})_2$	2	737.1
Magnesium Aluminum Phosphide	$\text{Mg}_3\text{AlP}_3 + 9\text{H}_2\text{O} \rightarrow 3\text{PH}_3 + \text{Al}(\text{OH})_3 + 3\text{Mg}(\text{OH})_2$	3	1865.2

Summary of Animal Data Relevant to AEGL-1

No animal data are available for the derivation of AEGL-1 for the metal phosphides considered in this appendix.

Derivation of AEGL-1

No human or animal data are consistent with the effects defined by AEGL-1. Data were also insufficient for derivation of AEGL-1 values for phosphine; thus phosphine cannot be used as a surrogate. Therefore, AEGL-1 values for the metal phosphides considered in this appendix are not recommended (Table D-11).

D.IV. RATIONALE AND AEGL-2

Summary of Human Data Relevant to AEGL-2

No human data are available for the derivation of AEGL-2 for the metal phosphides considered in this appendix.

TABLE D-11 AEGL-1 Values for Metal Phosphides

Compound	10 min	30 min	1 h	4 h	8 h
Aluminum Phosphide	NR	NR	NR	NR	NR
Potassium Phosphide	NR	NR	NR	NR	NR
Sodium Phosphide	NR	NR	NR	NR	NR
Zinc Phosphide	NR	NR	NR	NR	NR
Calcium Phosphide	NR	NR	NR	NR	NR
Magnesium Phosphide	NR	NR	NR	NR	NR
Strontium Phosphide	NR	NR	NR	NR	NR
Magnesium Aluminum Phosphide	NR	NR	NR	NR	NR

NR: not recommended. Absence of an AEGL-1 does not imply that exposure below the AEGL-2 is without adverse effects.

Summary of Animal Data Relevant to AEGL-2

No animal data are available for the derivation of AEGL-2 for the metal phosphides considered in this appendix.

Derivation of AEGL-2

In the absence of appropriate chemical-specific data for the metal phosphides considered in this appendix, the AEGL-2 values for phosphine will be used to obtain AEGL-2 values for the metal phosphides. The use of phosphine as a surrogate for the metal phosphides is deemed appropriate because qualitative (clinical signs) and quantitative (phosphine blood level) data suggest that the phosphine hydrolysis product is responsible for acute toxicity from metal phosphides. The phosphine AEGL-2 values will be used as target values for calculating the concentrations of metal phosphide needed to generate the phosphine AEGL values. Calculations were done using the methodology in NRC (2001) and are for 25 degrees C and 760 mm Hg. The metal phosphide values for AEGL-2 are given in Table D-12.

D.V. RATIONALE AND AEGL-3

Summary of Human Data Relevant to AEGL-3

No human data are available for the derivation of AEGL-3 for the metal phosphides considered in this appendix.

Summary of Animal Data Relevant to AEGL-3

No animal data are available for the derivation of AEGL-3 for the metal phosphides considered in this appendix.

TABLE D-12 AEGL-2 Values For Metal Phosphides^a

Compound	10-min	30-min	1-hr	4-hr	8-hr
Aluminum Phosphide	9.5 mg/m ³	9.5 mg/m ³	4.7 mg/m ³	1.2 mg/m ³	0.59 mg/m ³
Potassium Phosphide	24 mg/m ³	24 mg/m ³	12 mg/m ³	3.0 mg/m ³	1.5 mg/m ³
Sodium Phosphide	16 mg/m ³	16 mg/m ³	8.2 mg/m ³	2.0 mg/m ³	1.0 mg/m ³
Zinc Phosphide	21 mg/m ³	21 mg/m ³	11 mg/m ³	2.6 mg/m ³	1.3 mg/m ³
Calcium Phosphide	15 mg/m ³	15 mg/m ³	7.4 mg/m ³	1.9 mg/m ³	0.93 mg/m ³
Magnesium Phosphide	11 mg/m ³	11 mg/m ³	5.5 mg/m ³	1.4 mg/m ³	0.69 mg/m ³
Strontium Phosphide	27 mg/m ³	27 mg/m ³	13 mg/m ³	3.3 mg/m ³	1.7 mg/m ³
Magnesium Aluminum Phosphide	11 mg/m ³	11 mg/m ³	5.3 mg/m ³	1.3 mg/m ³	0.66 mg/m ³

^aThese airborne concentrations will produce the equivalent AEGL values for phosphine.

Derivation of AEGL-3

In the absence of appropriate chemical-specific data for the metal phosphides considered in this appendix, the AEGL-3 values for phosphine will be used to obtain AEGL-3 values for the metal phosphides. The use of phosphine as a surrogate for the metal phosphides is deemed appropriate because qualitative (clinical signs) and quantitative (phosphine blood level) data suggest that the phosphine hydrolysis product is responsible for acute toxicity from metal phosphides. The phosphine AEGL-3 values will be used as target values for calculating the concentrations of metal phosphide needed to generate the phosphine AEGL values. Calculations were done using the methodology in NRC (2001) and are for 25 degrees C and 760 mm Hg. The metal phosphide values for AEGL-3 are given in Table D-13.

D.VI. Comparison with Other Standards and Criteria

No other exposure criteria or guidelines were located for the metal phosphides.

TABLE D-13 AEGL-3 Values for Metal Phosphides^a

Compound	10-min	30-min	1-hr	4-hr	8-hr
Aluminum Phosphide	17 mg/m ³	17 mg/m ³	8.5 mg/m ³	2.1 mg/m ³	1.1 mg/m ³
Potassium Phosphide	44 mg/m ³	44 mg/m ³	22 mg/m ³	5.5 mg/m ³	2.7 mg/m ³
Sodium Phosphide	29 mg/m ³	29 mg/m ³	15 mg/m ³	3.7 mg/m ³	1.8 mg/m ³
Zinc Phosphide	38 mg/m ³	38 mg/m ³	19 mg/m ³	4.8 mg/m ³	2.4 mg/m ³
Calcium Phosphide	27 mg/m ³	27 mg/m ³	13 mg/m ³	3.4 mg/m ³	1.7 mg/m ³
Magnesium Phosphide	20 mg/m ³	20 mg/m ³	9.9 mg/m ³	2.5 mg/m ³	1.2 mg/m ³
Strontium Phosphide	48 mg/m ³	48 mg/m ³	24 mg/m ³	6.0 mg/m ³	3.0 mg/m ³
Magnesium Aluminum Phosphide	19 mg/m ³	19 mg/m ³	9.5 mg/m ³	2.4 mg/m ³	1.2 mg/m ³

^aThese airborne concentrations will produce the equivalent AEGL values for phosphine.

 注：本物質の特性理解のため、参考として急性曝露ガイドライン濃度 (AEGL)の原文のURLを記載する。

日本語ICSC

なし

AEGL (原文)

https://www.epa.gov/sites/default/files/2014-11/documents/phosphine_metal_phosphides_final_volume6_2007_5.pdf